



Nuclear Chemistry Cumulative Exam – Wednesday, Feb 16, 2011

This examination is concerned with nuclear reactions in nature as discussed in the textbook Modern Nuclear Chemistry by Loveland, Morrissey, and Seaborg. Possibly useful equations and constants are provided on page 4. There are a total of 100 points on this exam.

- (5 pts) Shown below is the chart of the nuclides with approximate paths for three different astrophysical processes. Stable nuclei are indicated by black squares. Match each numbered path with the name of the process responsible, choosing from either s-process (slow neutron capture process), rp-process (rapid-proton capture process), or r-process (rapid neutron capture process).



- (15 pts) Shown below is a schematic diagram of the relative solar abundances of the elements as a function of atomic mass. Heavy elements can be created in multiple processes. Two common ones are the slow-neutron capture process, characterized by neutron capture time scales that are long compared to beta decay lifetimes and the rapid-neutron capture process in which neutron capture time scales are shorter than beta decay lifetimes.




Identify the peaks labeled 1 and 2 as originating from either the s-process or r-process and explain why peak 1 is at a lower mass compared to peak 2.

3. (20 pts) The temperature averaged reaction rate per particle pair $\langle \sigma v \rangle$ is given by the equation



$$\langle \sigma v \rangle = \left(\frac{8}{\pi\mu}\right)^{1/2} \frac{1}{(kT)^{3/2}} \int_0^{\infty} S(E) \exp\left(-\frac{E}{kT} - \frac{b}{E^{1/2}}\right) dE$$

where μ is the reduced mass, k is Boltzmann's constant, T is the gas temperature, $S(E)$ is the astrophysical S-factor, and b is $0.989Z_1Z_2\mu^{1/2}(\text{MeV})^{1/2}$. The integral is dominated by the exponential term and represents the overlap between the Maxwell-Boltzmann distribution and a sub-barrier tunneling probability called the Gamow factor. For the following questions consider the simple non-resonant fusion reaction between two protons in a stellar environment.

- Estimate the Coulomb barrier for this reaction
- Reproduce the following figure in your exam book and qualitatively sketch the Maxwell-Boltzmann distribution and Gamow factor. Based on your sketch, indicate the location of the Gamow peak and explain its significance?
- Show that the maximum of the Gamow peak occurs at $E_0 = (bkT/2)^{2/3}$




4. (20 pts) Consider the sequence of three stable Te isotopes; ^{123}Te , ^{124}Te , and ^{125}Te . ^{123}Te and ^{124}Te are only produced from the s-process while the abundance of ^{125}Te is produced through both s- and r-processes. For the following questions assume that the neutron capture cross section is 808 mb, 155 mb, and 431 mb for ^{123}Te , ^{124}Te , and ^{125}Te , respectively and the abundance of ^{124}Te and ^{125}Te are 0.2319 and 0.3437 per 10^6 Si atoms, respectively.
- What is the abundance of ^{123}Te in the s-process at equilibrium?
 - What percentage of ^{125}Te is produced by the r-process?
5. (10 pts) There are two different reaction mechanisms for the fusion of two protons into a deuterium nucleus:



Explain why reaction (1) results in a broad spectrum of neutrino energies and why reaction (2) is a monoenergetic neutrino source.

6. (10 pts) The binding energy per nucleon as a function of mass number is shown below. Based on the figure, explain why fusion processes within a star will not contribute to the creation of elements with masses heavier than approximately $A \sim 60$.





7. (10 pts) The energy produced by the sun is approximately $4 \cdot 10^{26}$ J/s. Assuming that the energy production is from hydrogen burning within the sun according to the overall reaction




Estimate the rate of hydrogen consumption in the sun in grams / second.

8. (10 pts) The National Ignition Facility is being constructed at Livermore National Laboratory to demonstrate energy gain - the generation of energy from a fusion reaction in excess of the energy required to initiate the reaction. With three isotopes of hydrogen available, H (hydrogen), D (deuterium), and T (tritium) there are multiple possible reactions such as:



The reaction rate as a function of temperature for a variety of fusion reactions is shown below.



Which of the three reactions, D+D, T+T, or D+T has the greatest chance of success at NIF based on both energy generation and reaction rate?

$$P(v) = \left(\frac{m}{2\pi kT}\right)^{3/2} \exp\left(\frac{-mv^2}{2kT}\right)$$

$$\sigma(E) = \frac{1}{E} \exp\left[-31.29Z_1Z_2\left(\frac{\mu}{E}\right)^{1/2}\right]$$

$$\sigma(E) = \pi\lambda^2 \left[\frac{2J_r + 1}{(2J_x + 1)(2J_y + 1)}\right] \frac{\Gamma_{in}\Gamma_{out}}{(E - E_r)^2 + \frac{\Gamma_{tot}^2}{4}}$$

$$R = N\sigma\phi$$

$$\frac{dN_a}{dt} = \sigma_{A-1}N_{A-1} - \sigma_A N_A$$

$$Q = \sum \Delta(\text{reactants}) - \sum \Delta(\text{products})$$

$$T(K) = \frac{1.5 \times 10^{10}}{\sqrt{t(s)}}$$

$$V_C = \frac{z_1 z_2 e^2}{r_1 + r_2}$$

Table of Mass Excess, Δ

Nuclide	Δ (MeV)	Nuclide	Δ (MeV)
n	8.071	⁵ Li	11.68
¹ H	7.289	⁶ Li	14.087
² H	13.136	⁷ Li	14.908
³ H	14.590	⁸ Li	20.947
³ He	14.931	⁷ Be	15.770
⁴ He	2.425	⁸ Be	4.942
⁵ He	11.39	⁹ Be	11.348
⁶ He	17.595	¹⁰ Be	12.607

Constants:

Atomic mass unit	931.494 MeV	Atomic mass unit	1.66054*10 ⁻²⁷ kg
Proton radius	1.3214*10 ⁻¹⁵ m	Neutron radius	1.3196*10 ⁻¹⁵ m
m _e	0.510999 MeV	m _e	9.10939*10 ⁻²⁷ kg
m _p	938.272 MeV	m _p	1.67262*10 ⁻²⁷ kg
m _n	939.566 MeV	m _n	1.67493*10 ⁻²⁷ kg
c	2.99792458*10 ⁸ m/s	N _a	6.022*10 ²³ mol ⁻¹
h	6.62607*10 ⁻³⁴ J/s	k	1.3806*10 ⁻²³ J/K
e ² /hc	137.036	hc	197.327 MeV fm

Conversions:

$$1 \text{ eV} = 1.602177 \times 10^{-19} \text{ J}$$

$$1 \text{ Ci} = 3.700 \times 10^{10} \text{ Bq}$$

$$1 \text{ erg} = 10^{-7} \text{ J}$$