



NAME _____


PID _____

1) Calculate the degree of unsaturation for the formula (C₉H₂₀) (1 pt)¹H-NMR (Show structure fragment work in spectrum) (2 pts)¹³C-NMR (Show structure fragments work in spectrum) (2 pts).

Sometimes the DEPT function stops working or the undergraduate NMRs at your University don't have the DEPT option. This is the situation here.



2) Structure here. (2 pts)



Structure MUST BE PLACED Here
(show work for credit)


NAME _____

PID _____


3) Calculate the degree of unsaturation for the formula (C₅H₁₂O) 0 (1 pt)

NO OH
O RDB


¹H-NMR (Show structure fragments work in spectrum) (2 pts)



¹³C-NMR (Show structure fragments work in spectrum) (2 pts)



4) Structure here. (2 pts)





Structure MUST BE PLACED Here
(show work for credit)



5) Calculate the degree of unsaturation for the formula (C₅H₁₂O) 1 (1 pt)

C₇H₁₄O


¹H-NMR (Show structure fragments work in spectrum) (2 pts)




¹³C-NMR (Show structure fragments work in spectrum) (2 pts)



6) Structure here (2 pts)



Structure MUST BE PLACED Here (show work for credit)




NAME _____

PID _____

7) Formula (C_5H_9BrO)


① RDB

1H -NMR (Show structure fragments work in spectrum) (2 pts)




^{13}C -NMR (Show structure fragments work in spectrum) (2 pts).

Sometimes the DEPT function stops working or the undergraduate NMRs at your University don't have the DEPT option. This is the situation here.




8) Structure here (2 pts)




Structure MUST BE PLACED Here
(show work for credit)


9) Identify the indicated sets of protons as (U)nrelated, (H)omotopic, (E)nantiotopic or (D)iastereotopic (3 pts)




10) How many kinds of nonequivalent carbons (C) and protons (H) are present in the following? That is, how many unique ¹H and ¹³C signals will you observed in an NMR spectrum for each. (4 pts)



11) Arrange each set from 1 (slowest) to 3 (fastest) reactant in an S_N2 reaction. (3 pts)




12) Draw the compound below with wedged (—) and/or dashed (-----) lines to show the 3-D aspect of the chirality centers below. (3 pts)




NAME _____

PID _____


13) Identify whether the following molecules are (E)nantimers, (I)dentical, (D)iastereoisomers, (C)onstitutional isomers, or (N)ot isomers. Place appropriate letter in the box. (2 pts)



14) Assign R or S configuration to the chiral center(s). (5 pts)




15) Name. (3 pts) Use stereochemical descriptors with their numerical location(s) in the name.



(1R, 2S)-1-chloro-3-propylcyclohexane

16) (7 pts total) Under the condition shown, draw ALL the product(s) for the MOST LIKELY reaction mechanism below (Include stereochemistry, if necessary). (6 pts) Circle the reaction mechanism below the arrow. (1 pts) Take your time.




Draw HMPA (2 pt Bonus)


NAME _____

PID _____

17) (7 pts total) Under the condition shown, draw ALL the product(s) for the MOST LIKELY reaction mechanism below (Include stereochemistry, if necessary). (6 pts) Circle the reaction mechanism below the arrow. (1 pts). Note the reactant above arrow; remember what it can ONLY function as. Take your time.




18) Draw. (3 pts)




3-isopropyl-2,4-dimethyl-2-pentene

19) Name. Include E/Z stereochemistry in name, if necessary. (3 pts)



* both accepted.

20) (9 pts total) Assign E or Z configuration (1 pt). CIRCLE both highest priority groups for in each. (1 pt each circle)




4213



NAME _____

PID _____

21) Arrange the following in order of increasing stability (1 is LEAST stable; 4 is MOST stable).




22) Draw the MAJOR (most abundant) product made in the reaction. IF NEW STEREOCENTER(S) IS/ARE MADE, PLACE A DOT (•) ON THE CHIRALITY CENTER(S). (6 pts)




23) Identify the products of the following Elimination reactions. (6 pts)

- 1) If only a MAJOR product is made, draw an X in the MINOR box.
- 2) If no reaction draw X in both boxes
- 3) Disregard trans/cis stereochemistry as MAJOR/MINOR for this problem.




24) (7 pts total) Identify the STEREOISOMERIC products of the following Elimination reactions (6 pts). Circle the reaction mechanism below the arrow (1 pt each). If TWO PRODUCTS ARE NOT MADE, THE DRAW AN X IN THE BOX DESIGNATED 'MINOR'



Bonus: Must be completely correct for credit 5 points

Place numbers 1, 2, 3, 4, 5 in the BOXES to designate C1, C2, C3, C4, C5, and write R or S stereochemical descriptors in the CIRCLES of pentamethylcyclohexane below



sets priority over "S"
∴ this in C₁

